

DM554  
Linear and Integer Programming

Lecture 3  
**Matrices and Vectors**

Marco Chiarandini

Department of Mathematics & Computer Science  
University of Southern Denmark


1. Geometric Insight

2. Linear Systems



1. Geometric Insight

2. Linear Systems

- Set  $\mathbb{R}$  can be represented by **real-number line**. Set  $\mathbb{R}^2$  of real number pairs  $(a_1, a_2)$  can be represented by the **Cartesian plane**.
- To a point in the plane  $A = (a_1, a_2)$  it is associated a **position vector**  $\mathbf{a} = (a_1, a_2)^T$ , representing the displacement from the origin  $(0, 0)$ .  $\diamond$



- Two displacement vectors of same **length** and **direction** are considered to be equal even if they do not both start from the origin
- If object displaced from O to P by displacement **p** and from P to Q by displacement **v**, then the total displacement satisfies **q = p + v = v + q**



- **v = q - p**, think of **v** as the vector that is added to **p** to obtain **q**.

- the **length** of a vector  $\mathbf{a} = (a_1, a_2)^T$  is denoted by  $\|\mathbf{a}\|$  and from Pythagoras

$$\|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = \sqrt{\langle \mathbf{a}, \mathbf{a} \rangle}$$


- the **direction** is given by the components of the vector
- the unit vector can be derived from

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v}$$

### Theorem (Inner Product)


Let  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^2$  and let  $\theta$  denote the angle between them. Then,

$$\langle \mathbf{a}, \mathbf{b} \rangle = \|\mathbf{a}\| \|\mathbf{b}\| \cos \theta$$



Two vectors  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal (or normal or perpendicular) if and only if  $\langle \mathbf{a}, \mathbf{b} \rangle = 0$ .

# Vectors in $\mathbb{R}^3$



$$\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}$$

$$\|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

$$\langle \mathbf{a}, \mathbf{b} \rangle = \|\mathbf{a}\| \|\mathbf{b}\| \cos \theta$$

# Lines in $\mathbb{R}^2$

- Cartesian line equation  $y = ax + b$

- another way is by giving position vectors.

We can let  $x = t$  where  $t$  is any real number. Then

$y = ax + b = at + b$ . Hence the position vector  $\mathbf{x} = (x, y)^T$

$$\mathbf{x} = \begin{bmatrix} t \\ at + b \end{bmatrix} = t \begin{bmatrix} 1 \\ a \end{bmatrix} + \begin{bmatrix} 0 \\ b \end{bmatrix} = t\mathbf{v} + (0, b)^T, \quad t \in \mathbb{R}$$


- To derive the Cartesian equation: locate one particular point on the line, eg, the  $y$  intercept. Then the position vector of any point on the line is a sum of two displacements, first going to the point and then along the direction of the line. Try with  $P = (-1, 1)$  and  $Q = (3, 2)$
- In general, any line in  $\mathbb{R}^2$  is given by a vector equation with one parameter of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}$$

where  $\mathbf{x}$  is the position vector,  $\mathbf{p}$  is any particular point and  $\mathbf{v}$  is the direction of the line



# Lines in $\mathbb{R}^3$



$$\mathbf{x} = \mathbf{p} + t\mathbf{v}$$

$$\mathbf{x} = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$\mathbf{x} = \begin{bmatrix} 3 \\ 7 \\ 2 \end{bmatrix} + s \begin{bmatrix} -3 \\ -6 \\ 3 \end{bmatrix}, \quad s, t \in \mathbb{R}$$

Are these lines intersecting?  
What is the Cartesian equation of the first?

In  $\mathbb{R}^2$ , two lines are:

- parallel
- intersecting in a unique point

In  $\mathbb{R}^3$ , two lines are:

- parallel
- intersecting in a unique point
- skew (lay on two parallel planes)

What about these lines? Do they intersect? Are they coplanar?

$$L_1 : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$L_2 : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \\ 1 \end{bmatrix} + t \begin{bmatrix} -2 \\ 1 \\ 7 \end{bmatrix}$$

Vector parametric equation:


- The position of vectors of points on a plane is described by:

$$\mathbf{x} = \mathbf{p} + s\mathbf{v} + t\mathbf{w}, \quad s, t \in \mathbb{R}$$

provided  $\mathbf{v}$  and  $\mathbf{w}$  are non-zero and not parallel.

( $\mathbf{p}$  position vector,  $\mathbf{v}$  and  $\mathbf{w}$  displacement vectors).

- How is the plane through the origin? What if  $\mathbf{v}$  and  $\mathbf{w}$  are parallel?
- Two intersecting lines determine a plane. What is its description?



Cartesian equation:


- Let  $\mathbf{n}$  be a given vector in  $\mathbb{R}^3$ . All positions represented by position vectors  $\mathbf{x}$  that are orthogonal to  $\mathbf{n}$  describe a plane through the origin. ( $\mathbf{n}$  is called a normal vector to the plane)
- Vectors  $\mathbf{n}$  and  $\mathbf{x}$  are orthogonal iff

$$\langle \mathbf{n}, \mathbf{x} \rangle = 0,$$

hence this equation describes a plane.

If  $\mathbf{n} = (a, b, c)^T$  and  $\mathbf{x} = (x, y, z)^T$ , then the equation becomes:

$$ax + by + cz = 0$$



- For a point  $P$  on the plane with position vector  $\mathbf{p}$  and a position vector  $\mathbf{x}$  of any other point on the plane, the displacement vector  $\mathbf{x} - \mathbf{p}$  lies on the plane and  $\mathbf{n} \perp \mathbf{x} - \mathbf{p}$
- Conversely, if the position vector  $\mathbf{x}$  of a point is such that

$$\langle \mathbf{n}, \mathbf{x} - \mathbf{p} \rangle = 0$$

then the point represented by  $\mathbf{x}$  lies on the plane.

- hence,  $\langle \mathbf{n}, \mathbf{x} \rangle = \langle \mathbf{n}, \mathbf{p} \rangle = d$  and the equation becomes:

$$ax + by + cz = d$$

Eg.:  $2x - 3y - 5z = 2$  has  $\mathbf{n} = (2, -3, -5)^T$  and passes through  $(0, 0, e)$

Vector parametric equation  $\iff$  Cartesian equation

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = s \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + t \begin{bmatrix} 2 \\ 1 \\ 7 \end{bmatrix} = s\mathbf{v} + t\mathbf{w}, \quad s, t \in \mathbb{R}$$

$$3x - y + z = 0, \quad \mathbf{n} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\langle \mathbf{n}, \mathbf{v} \rangle = 0, \langle \mathbf{n}, \mathbf{w} \rangle = 0 \text{ and } \langle \mathbf{n}, s\mathbf{v} + t\mathbf{w} \rangle = 0 \text{ for } s, t \in \mathbb{R}$$

What changes if the plane does not pass through the origin?



Are the two following planes parallel?

$$x + 2y - 3z = 0 \text{ and } -2x - 4y + 6z = 4$$

and these?

$$x + 2y - 3z = 0 \text{ and } x - 2y + 5z = 4$$

# Lines and Hyperplanes in $\mathbb{R}^n$

- Point in  $\mathbb{R}^n$ :  $\mathbf{a} = (a_1, a_2, \dots, a_n)^T$
- Length of a vector  $\mathbf{x} = (x_1, x_2, \dots, x_n)^T$

$$\|\mathbf{x}\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}.$$

- The vectors in  $\mathbb{R}^n$  are orthogonal iff

$$\langle \mathbf{v}, \mathbf{w} \rangle = 0.$$

- Line:

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}, \quad t \in \mathbb{R}$$

How many Cartesian equations?

- The set of points  $(x_1, x_2, \dots, x_n)$  that satisfy a Cartesian equation

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = d$$

is called **hyperplane**. ( $\langle \mathbf{n}, \mathbf{x} - \mathbf{p} \rangle = 0$ .) What is the vector equation?

1. Geometric Insight

2. Linear Systems

# Systems of Linear Equations

Definition (System of linear equations, aka linear system)

A system of  $m$  linear equations in  $n$  unknowns  $x_1, x_2, \dots, x_n$  is a set of  $m$  equations of the form

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m \end{aligned}$$

The numbers  $a_{ij}$  are known as the **coefficients** of the system.

We say that  $s_1, s_2, \dots, s_n$  is a **solution** of the system if all  $m$  equations hold true when

$$x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$$

# Examples

$$\begin{aligned}x_1 + x_2 + x_3 + x_4 + x_5 &= 3 \\2x_1 + x_2 + x_3 + x_4 + 2x_5 &= 4 \\x_1 - x_2 - x_3 + x_4 + x_5 &= 5 \\x_1 &+ x_4 + x_5 = 4\end{aligned}$$

has solution

$$x_1 = -1, x_2 = -2, x_3 = 1, x_4 = 3, x_5 = 2.$$

Is it the only one?

$$\begin{aligned}x_1 + x_2 + x_3 + x_4 + x_5 &= 3 \\2x_1 + x_2 + x_3 + x_4 + 2x_5 &= 4 \\x_1 - x_2 - x_3 + x_4 + x_5 &= 5 \\x_1 &+ x_4 + x_5 = 6\end{aligned}$$

has no solutions

## Definition (Coefficient Matrix)

The matrix  $A = (a_{ij})$ , whose  $(i, j)$  entry is the coefficient  $a_{ij}$  of the system of linear equations is called the **coefficient matrix**.

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

Let  $\mathbf{x} = (x_1, x_2, \dots, x_n)^T$  then

$$\begin{matrix} & m \times n & & n \times 1 & & n \times 1 \\ \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} & & \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} & = & \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots & \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix} \end{matrix}$$

hence, the linear system can be written also as  $A\mathbf{x} = \mathbf{b}$

# Row operations

How do we find solutions?

$$\begin{array}{l} \text{I:} \\ \text{II:} \\ \text{III:} \end{array} \left| \begin{array}{l} x_1 + x_2 + x_3 = 3 \\ 2x_1 + x_2 + x_3 = 4 \\ x_1 - x_2 + 2x_3 = 5 \end{array} \right.$$

Eliminate one of the variables from two of the equations

$$\begin{array}{l} \text{I}'=\text{I:} \\ \text{II}'=\text{II}-2*\text{I:} \\ \text{III}'=\text{III:} \end{array} \left| \begin{array}{l} x_1 + x_2 + x_3 = 3 \\ -x_2 - x_3 = -2 \\ x_1 - x_2 + 2x_3 = 5 \end{array} \right.$$

$$\begin{array}{l} \text{I}'=\text{I}': \\ \text{II}'=\text{II}': \\ \text{III}'=\text{III}'-\text{I}: \end{array} \left| \begin{array}{l} x_1 + x_2 + x_3 = 3 \\ -x_2 - x_3 = -2 \\ x_1 - x_2 + 2x_3 = 5 \end{array} \right.$$

We can now eliminate one of the variables in the last two equations to obtain the solution

Row operations that do not alter solutions:

O1: multiply both sides of an equation by a non-zero constant

O2: interchange two equations

O3: add a multiple of one equation to another

These operations only act on the coefficients of the system

For a system  $A\mathbf{x} = \mathbf{b}$ :

$$[A|\mathbf{b}] = \begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & 1 & 1 & 4 \\ 1 & -1 & 2 & 5 \end{bmatrix}$$



# Augmented Matrix

## Definition (Augmented Matrix and Elementary row operations)

For a system of linear equations  $A\mathbf{x} = \mathbf{b}$  with

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

the augmented matrix of the system and the row operations are:

$$[A | \mathbf{b}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix}$$

RO1: multiply a row by a non-zero constant

RO2: interchange two rows

RO3: add a multiple of one row to another