

DM545  
Linear and Integer Programming

Lecture 6  
More on Duality

Marco Chiarandini

Department of Mathematics & Computer Science  
University of Southern Denmark

## 1. Derivation

Geometric Interpretation

Lagrangian Duality

Dual Simplex

## 2. Sensitivity Analysis

- Derivation:
  1. economic interpretation
  2. bounding
  3. multipliers
  4. recipe
  5. Lagrangian
- Theory:
  - Symmetry
  - Weak duality theorem
  - Strong duality theorem
  - Complementary slackness theorem
- Dual Simplex
- Sensitivity Analysis, Economic interpretation

## 1. Derivation

Geometric Interpretation

Lagrangian Duality

Dual Simplex

## 2. Sensitivity Analysis

Dual variables  $\mathbf{y}$  in one-to-one correspondence with the constraints:

Primal problem:

$$\begin{aligned} \max \quad & z = \mathbf{c}^T \mathbf{x} \\ & A\mathbf{x} = \mathbf{b} \\ & \mathbf{x} \geq 0 \end{aligned}$$

Dual Problem:

$$\begin{aligned} \min \quad & w = \mathbf{b}^T \mathbf{y} \\ & A^T \mathbf{y} \geq \mathbf{c} \\ & \mathbf{y} \in \mathbb{R}^m \end{aligned}$$

- Basic feasible solutions of (P) give immediate lower bounds on the optimal value  $z^*$ . Is there a simple way to get upper bounds?
- The optimal solution must satisfy any linear combination  $\mathbf{y} \in \mathbb{R}^m$  of the equality constraints.
- If we can construct a linear combination of the equality constraints  $\mathbf{y}^T(A\mathbf{x}) = \mathbf{y}^T\mathbf{b}$ , for  $\mathbf{y} \in \mathbb{R}^m$ , such that  $\mathbf{c}^T\mathbf{x} \leq \mathbf{y}^T(A\mathbf{x})$ , then  $\mathbf{y}^T(A\mathbf{x}) = \mathbf{y}^T\mathbf{b}$  is an upper bound on  $z^*$ .

## 1. Derivation

Geometric Interpretation


Lagrangian Duality

Dual Simplex

## 2. Sensitivity Analysis

# Geometric Interpretation

$$\begin{aligned} \max \quad & x_1 + x_2 = z \\ & 2x_1 + x_2 \leq 14 \\ & -x_1 + 2x_2 \leq 8 \\ & 2x_1 - x_2 \leq 10 \\ & x_1, x_2 \geq 0 \end{aligned}$$




Feasible sol  $x^* = (4, 6)$  yields  $z^* = 10$ . To prove that it is optimal we need to verify that  $y^* = (3/5, 1/5, 0)$  is a feasible solution of  $D$ :

$$\begin{aligned} \min \quad & 14y_1 + 8y_2 + 10y_3 = w \\ & 2y_1 - y_2 + 2y_3 \geq 1 \\ & y_1 + 2y_2 - y_3 \geq 1 \\ & y_1, y_2, y_3 \geq 0 \end{aligned}$$

and that  $w^* = 10$

$$\frac{\frac{3}{5} \cdot (2x_1 + x_2 \leq 14) + \frac{1}{5} \cdot (-x_1 + 2x_2 \leq 8)}{x_1 + x_2 \leq 10}$$



$$(2v - w)x_1 + (v + 2w)x_2 \leq 14v + 8w$$

set of halfplanes that contain the feasibility region of  $P$  and pass through  $[4, 6]$

$$2v - w \geq 1$$


$$v + 2w \geq 1$$

Example of boundary lines among those allowed:

$$v = 1, w = 0 \implies 2x_1 + x_2 = 14$$

$$v = 1, w = 1 \implies x_1 + 3x_2 = 22$$

$$v = 2, w = 1 \implies 3x_1 + 4x_2 = 36$$





## 1. Derivation

Geometric Interpretation

**Lagrangian Duality**

Dual Simplex

## 2. Sensitivity Analysis

# Lagrangian Duality

**Relaxation:** if a problem is hard to solve then find an easier problem resembling the original one that provides information in terms of bounds. Then, search for the strongest bounds.

$$\begin{aligned} \min \quad & 13x_1 + 6x_2 + 4x_3 + 12x_4 \\ & 2x_1 + 3x_2 + 4x_3 + 5x_4 = 7 \\ & 3x_1 + \quad + 2x_3 + 4x_4 = 2 \\ & x_1, x_2, x_3, x_4 \geq 0 \end{aligned}$$

We wish to reduce to a problem easier to solve, ie:

$$\begin{aligned} \min \quad & c_1x_1 + c_2x_2 + \dots + c_nx_n \\ & x_1, x_2, \dots, x_n \geq 0 \end{aligned}$$

solvable by inspection: if  $c < 0$  then  $x = +\infty$ , if  $c \geq 0$  then  $x = 0$ .

measure of violation of the constraints:

$$\begin{aligned} & 7 - (2x_1 + 3x_2 + 4x_3 + 5x_4) \\ & 2 - (3x_1 + \quad + 2x_3 + 4x_4) \end{aligned}$$

We relax these measures in obj. function with Lagrangian multipliers  $y_1, y_2$ .  
We obtain a family of problems:

$$PR(y_1, y_2) = \min_{x_1, x_2, x_3, x_4 \geq 0} \left\{ \begin{array}{l} 13x_1 + 6x_2 + 4x_3 + 12x_4 \\ +y_1(7 - 2x_1 - 3x_2 - 4x_3 - 5x_4) \\ +y_2(2 - 3x_1 - 2x_3 - 4x_4) \end{array} \right\}$$

1. for all  $y_1, y_2 \in \mathbb{R} : \text{opt}(PR(y_1, y_2)) \leq \text{opt}(P)$
2.  $\max_{y_1, y_2 \in \mathbb{R}} \{\text{opt}(PR(y_1, y_2))\} \leq \text{opt}(P)$

PR is easy to solve.

(It can be also seen as a proof of the weak duality theorem)

$$PR(y_1, y_2) = \min_{x_1, x_2, x_3, x_4 \geq 0} \left\{ \begin{array}{l} (13 - 2y_2 - 3y_1) x_1 \\ + (6 - 3y_1) x_2 \\ + (4 - 2y_2) x_3 \\ + (12 - 5y_1 - 4y_2) x_4 \\ + 7y_1 + 2y_2 \end{array} \right\}$$

if coeff. of  $x$  is  $< 0$  then bound is  $-\infty$  then LB is useless

$$(13 - 2y_2 - 3y_1) \geq 0$$

$$(6 - 3y_1) \geq 0$$

$$(4 - 2y_2) \geq 0$$

$$(12 - 5y_1 - 4y_2) \geq 0$$

If they all hold then we are left with  $7y_1 + 2y_2$  because all go to 0.

$$\max 7y_1 + 2y_2$$

$$2y_2 + 3y_1 \leq 13$$

$$3y_1 \leq 6$$

$$+ 2y_2 \leq 4$$

$$5y_1 + 4y_2 \leq 12$$

# General Formulation

$$\begin{array}{ll} \min & z = \mathbf{c}^T \mathbf{x} & \mathbf{c} \in \mathbb{R}^n \\ & \mathbf{A}\mathbf{x} = \mathbf{b} & \mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m \\ & \mathbf{x} \geq \mathbf{0} & \mathbf{x} \in \mathbb{R}^n \end{array}$$

$$\max_{\mathbf{y} \in \mathbb{R}^m} \left\{ \min_{\mathbf{x} \in \mathbb{R}_+^n} \{ \mathbf{c}^T \mathbf{x} + \mathbf{y}^T (\mathbf{b} - \mathbf{A}\mathbf{x}) \} \right\}$$

$$\max_{\mathbf{y} \in \mathbb{R}^m} \left\{ \min_{\mathbf{x} \in \mathbb{R}_+^n} \{ (\mathbf{c}^T - \mathbf{y}^T \mathbf{A})\mathbf{x} + \mathbf{y}^T \mathbf{b} \} \right\}$$

$$\begin{array}{ll} \max & \mathbf{b}^T \mathbf{y} \\ & \mathbf{A}^T \mathbf{y} \leq \mathbf{c} \\ & \mathbf{y} \in \mathbb{R}^m \end{array}$$

## 1. Derivation

Geometric Interpretation

Lagrangian Duality

**Dual Simplex**

## 2. Sensitivity Analysis

# Dual Simplex

- Dual simplex (Lemke, 1954): apply the simplex method to the dual problem and observe what happens in the primal tableau:

$$\begin{aligned} \max\{c^T x \mid Ax \leq b, x \geq 0\} &= \min\{b^T y \mid A^T y \geq c^T, y \geq 0\} \\ &= -\max\{-b^T y \mid -A^T x \leq -c^T, y \geq 0\} \end{aligned}$$

- We obtain a new algorithm for the primal problem: the dual simplex  
It corresponds to the primal simplex applied to the dual

Primal simplex on primal problem:

- pivot  $> 0$
- col  $c_j$  with wrong sign
- row:  
 $\min \left\{ \frac{b_i}{a_{ij}} : a_{ij} > 0, i = 1, \dots, m \right\}$

Dual simplex on primal problem:

- pivot  $< 0$
- row  $b_i < 0$   
(condition of feasibility)
- col:  
 $\min \left\{ \left| \frac{c_j}{a_{ij}} \right| : a_{ij} < 0, j = 1, 2, \dots, n + m \right\}$   
(least worsening solution)

0. (primal) simplex on primal problem (the one studied so far)
1. Now: dual simplex on primal problem  $\equiv$  primal simplex on dual problem (implemented as dual simplex, understood as primal simplex on dual problem)

Uses of 1.:

- The dual simplex can work better than the primal in some cases.  
Eg. since running time in practice between  $2m$  and  $3m$ , then if  $m = 99$  and  $n = 9$  then better the dual
- Infeasible start  
Dual based Phase I algorithm (Dual-primal algorithm)



# Dual Simplex for Phase I

## Example

Primal:

$$\begin{aligned}
 \max \quad & -x_1 - x_2 \\
 & -2x_1 - x_2 \leq 4 \\
 & -2x_1 + 4x_2 \leq -8 \\
 & -x_1 + 3x_2 \leq -7 \\
 & x_1, x_2 \geq 0
 \end{aligned}$$

Dual:

$$\begin{aligned}
 \min \quad & 4y_1 - 8y_2 - 7y_3 \\
 & -2y_1 - 2y_2 - y_3 \geq -1 \\
 & -y_1 + 4y_2 + 3y_3 \geq -1 \\
 & y_1, y_2, y_3 \geq 0
 \end{aligned}$$

- Initial tableau

	x1	x2	w1	w2	w3	-z	b
	-2	-1	1	0	0	0	4
	-2	4	0	1	0	0	-8
	-1	3	0	0	1	0	-7
	-1	-1	0	0	0	1	0

infeasible start

- $x_1$  enters,  $w_2$  leaves

- Initial tableau ( $\min by \equiv -\max -by$ )

	y1	y2	y3	z1	z2	-p	b
	2	2	1	1	0	0	1
	1	-4	-3	0	1	0	1
	-4	8	7	0	0	1	0

feasible start (thanks to  $-x_1 - x_2$ )

- $y_2$  enters,  $z_1$  leaves

- $x_1$  enters,  $w_2$  leaves

	$x_1$	$x_2$	$w_1$	$w_2$	$w_3$	$-z$	$b$
	0	-5	1	-1	0	0	12
	1	-2	0	-0.5	0	0	4
	0	1	0	-0.5	1	0	-3
	0	-3	0	-0.5	0	1	4

- $y_2$  enters,  $z_1$  leaves

	$y_1$	$y_2$	$y_3$	$z_1$	$z_2$	$-p$	$b$
	1	1	0.5	0.5	0	0	0.5
	5	0	-1	2	1	0	3
	-4	0	3	-12	0	1	-4

- $w_2$  enters,  $w_3$  leaves (note that we kept  $c_j < 0$ , ie, optimality)

	$x_1$	$x_2$	$w_1$	$w_2$	$w_3$	$-z$	$b$
	0	-7	1	0	-2	0	18
	1	-3	0	0	-1	0	7
	0	-2	0	1	-2	0	6
	0	-4	0	0	-1	1	7

- $y_3$  enters,  $y_2$  leaves

	$y_1$	$y_2$	$y_3$	$z_1$	$z_2$	$-p$	$b$
	2	2	1	1	0	0	1
	7	2	0	3	1	0	3
	-18	-6	0	-7	0	1	-7

- Derivation:
  1. bounding
  2. multipliers
  3. recipe
  4. Lagrangian
- Theory:
  - Symmetry
  - Weak duality theorem
  - Strong duality theorem
  - Complementary slackness theorem
- Dual Simplex
- Sensitivity Analysis, Economic interpretation

## 1. Derivation

Geometric Interpretation

Lagrangian Duality

Dual Simplex

## 2. Sensitivity Analysis

# Economic Interpretation

$$\begin{aligned} \max \quad & 5x_0 + 6x_1 + 8x_2 \\ & 6x_0 + 5x_1 + 10x_2 \leq 60 \\ & 8x_0 + 4x_1 + 4x_2 \leq 40 \\ & 4x_0 + 5x_1 + 6x_2 \leq 50 \\ & x_0, x_1, x_2 \geq 0 \end{aligned}$$

final tableau:

$x_0$	$x_1$	$x_2$	$s_1$	$s_2$	$s_3$	$-z$	$b$
	0	1		0			$5/2$
	1	0		0			7
	0	0		1			2
$-1/5$	0	0	$-1/5$	0	$-1$		$-62$

- Which are the values of variables, the reduced costs, the shadow prices (or marginal prices), the values of dual variables?
- If one slack variable  $> 0$  then overcapacity:  $s_2 = 2$  then the second constraint is not tight
- How many products can be produced at most? at most  $m$
- How much more expensive a product not selected should be?  
look at reduced costs:  $c_j - \pi a_j > 0$
- What is the value of extra capacity of manpower? In +1 out  $+1/5$

# Economic Interpretation

Game: Suppose two economic operators:

- P owns the factory and produces goods
- D is in the market buying and selling raw material and resources
- D asks P to close and sell him all resources
- P considers if the offer is convenient
- D wants to spend least possible
- $y$  are prices that D offers for the resources
- $\sum y_i b_i$  is the amount D has to pay to have all resources of P
- $\sum y_i a_{ij} \geq c_j$  total value to make  $j >$  price per unit of product
- P either sells all resources  $\sum y_i a_{ij}$  or produces product  $j$  ( $c_j$ )
- without  $\geq$  there would not be negotiation because P would be better off producing and selling
- ▶ at optimality the situation is indifferent (strong th.)
- ▶ resource 2 that was not totally utilized in the primal has been given value 0 in the dual. (complementary slackness th.) Plausible, since we do not use all the resource, likely to place not so much value on it.
- ▶ for product 0  $\sum y_i a_{ij} > c_j$  hence not profitable producing it. (complementary slackness th.)

# Sensitivity Analysis

aka Postoptimality Analysis

Derivation  
Sensitivity Analysis

Instead of solving each modified problem from scratch, exploit results obtained from solving the original problem.

$$\max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} = \mathbf{b}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\} \quad (*)$$

(I) changes to coefficients of objective function:

$$\max\{\tilde{\mathbf{c}}^T \mathbf{x} \mid A\mathbf{x} = \mathbf{b}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\} \quad (\text{primal})$$

$\mathbf{x}^*$  of (\*) remains feasible hence we can restart the simplex from  $\mathbf{x}^*$

(II) changes to RHS terms:  $\max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} = \tilde{\mathbf{b}}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\}$  (dual)  
 $\mathbf{x}^*$  optimal feasible solution of (\*)

basic sol  $\bar{\mathbf{x}}$  of (II):  $\bar{\mathbf{x}}_N = \mathbf{x}_N^*$ ,  $A_B \bar{\mathbf{x}}_B = \tilde{\mathbf{b}} - A_N \bar{\mathbf{x}}_N$

$\bar{\mathbf{x}}$  is dual feasible and we can start the dual simplex from there. If  $\tilde{\mathbf{b}}$  differs from  $\mathbf{b}$  only slightly it may be we are already optimal.

(III) introduce a new variable:

(primal)

$$\begin{aligned} \max \quad & \sum_{j=1}^6 c_j x_j \\ & \sum_{j=1}^6 a_{ij} x_j = b_i, \quad i = 1, \dots, 3 \\ & l_j \leq x_j \leq u_j, \quad j = 1, \dots, 6 \\ & [x_1^*, \dots, x_6^*] \text{ feasible} \end{aligned}$$

$$\begin{aligned} \max \quad & \sum_{j=1}^7 c_j x_j \\ & \sum_{j=1}^7 a_{ij} x_j = b_i, \quad i = 1, \dots, 3 \\ & l_j \leq x_j \leq u_j, \quad j = 1, \dots, 7 \\ & [x_1^*, \dots, x_6^*, 0] \text{ feasible} \end{aligned}$$

(IV) introduce a new constraint:

(dual)

$$\begin{aligned} & \sum_{j=1}^6 a_{4j} x_j = b_4 \\ & \sum_{j=1}^6 a_{5j} x_j = b_5 \\ & l_j \leq x_j \leq u_j \quad j = 7, 8 \end{aligned}$$

$$\begin{aligned} & [x_1^*, \dots, x_6^*] \text{ optimal} \\ & [x_1^*, \dots, x_6^*, x_7^*, x_8^*] \text{ feasible} \\ & x_7^* = b_4 - \sum_{j=1}^6 a_{4j} x_j^* \\ & x_8^* = b_5 - \sum_{j=1}^6 a_{5j} x_j^* \end{aligned}$$



# Examples

(I) Variation of reduced costs:

$$\begin{aligned} \max \quad & 6x_1 + 8x_2 \\ & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0 \end{aligned}$$

$$\begin{array}{c|cccccc} & x_1 & x_2 & x_3 & x_4 & -z & b \\ \hline x_3 & 5 & 10 & 1 & 0 & 0 & 60 \\ x_4 & 4 & 4 & 0 & 1 & 0 & 40 \\ \hline & 6 & 8 & 0 & 0 & 1 & 0 \end{array}$$

The last tableau gives the possibility to estimate the effect of variations

$$\begin{array}{c|cccccc} & x_1 & x_2 & x_3 & x_4 & -z & b \\ \hline x_2 & 0 & 1 & 1/5 & -1/4 & 0 & 2 \\ x_1 & 1 & 0 & -1/5 & 1/2 & 0 & 8 \\ \hline & 0 & 0 & -2/5 & -1 & 1 & -64 \end{array}$$

For a variable in basis the perturbation goes unchanged in the red. costs. Eg:

$$\max (6 + \delta)x_1 + 8x_2 \implies \bar{c}_1 = 1(6 + \delta) - \frac{2}{5} \cdot 5 - 1 \cdot 4 = \delta$$

then need to bring in canonical form and hence  $\delta$  changes the obj value.  
For a variable not in basis, if it changes the sign of the reduced cost  $\implies$   
worth bringing in basis  $\implies$  the  $\delta$  term propagates to other columns

## (II) Changes in RHS terms

$$\begin{array}{c|cccccc}
 & x_1 & x_2 & x_3 & x_4 & -z & b \\
 \hline
 x_3 & 5 & 10 & 1 & 0 & 0 & 60 + \delta \\
 x_4 & 4 & 4 & 0 & 1 & 0 & 40 + \epsilon \\
 \hline
 & 6 & 8 & 0 & 0 & 1 & 0
 \end{array}$$

$$\begin{array}{c|cccccc}
 & x_1 & x_2 & x_3 & x_4 & -z & b \\
 \hline
 x_2 & 0 & 1 & 1/5 & -1/4 & 0 & 2 + 1/5\delta - 1/4\epsilon \\
 x_1 & 1 & 0 & -1/5 & 1/2 & 0 & 8 - 1/5\delta + 1/2\epsilon \\
 \hline
 & 0 & 0 & -2/5 & -1 & 1 & -64 - 2/5\delta - \epsilon
 \end{array}$$

(It would be more convenient to augment the second. But let's take  $\epsilon = 0$ .)

If  $60 + \delta \implies$  all RHS terms change and we must check feasibility

Which are the multipliers for the first row?  $k_1 = \frac{1}{5}$ ,  $k_2 = -\frac{1}{4}$ ,  $k_3 = 0$


I:  $1/5(60 + \delta) - 1/4 \cdot 40 + 0 \cdot 0 = 12 + \delta/5 - 10 = 2 + \delta/5$

II:  $-1/5(60 + \delta) + 1/2 \cdot 40 + 0 \cdot 0 = -60/5 + 20 - \delta/5 = 8 - 1/5\delta$

Risk that RHS becomes negative

Eg: if  $\delta = -10 \implies$  tableau stays optimal but not feasible  $\implies$  apply dual simplex

# Graphical Representation



(III) Add a variable

$$\begin{aligned} \max \quad & 5x_0 + 6x_1 + 8x_2 \\ & 6x_0 + 5x_1 + 10x_2 \leq 60 \\ & 8x_0 + 4x_1 + 4x_2 \leq 40 \\ & x_0, x_1, x_2 \geq 0 \end{aligned}$$

Reduced cost of  $x_0$ ?  $c_j + \sum \pi_i a_{ij} = +1 \cdot 5 - \frac{2}{5} \cdot 6 + (-1)8 = -\frac{27}{5}$

To make worth entering in basis:

- increase its cost
- decrease the technological coefficient in constraint II:  
 $5 - 2/5 \cdot 6 - a_{20} > 0$

(IV) Add a constraint

$$\begin{aligned}
 \max \quad & 6x_1 + 8x_2 \\
 & 5x_1 + 10x_2 \leq 60 \\
 & 4x_1 + 4x_2 \leq 40 \\
 & 5x_1 + 6x_2 \leq 50 \\
 & x_1, x_2 \geq 0
 \end{aligned}$$

Final tableau not in canonical form, need to iterate with dual simplex

	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$-z$	$b$
$x_2$	0	1	$1/5$	$-1/4$		0	2
$x_1$	1	0	$-1/5$	$1/2$		0	8
	0	0	$5/5$	$6/4$	1	0	-2
	0	0	$-2/5$	-1	0	1	-64

(V) change in a technological coefficient:

$$\begin{array}{c|cccccc}
 & x_1 & x_2 & x_3 & x_4 & -z & b \\
 \hline
 x_3 & 5 & 10 + \delta & 1 & 0 & 0 & 60 \\
 x_4 & 4 & 4 & 0 & 1 & 0 & 40 \\
 \hline
 & 6 & 8 & 0 & 0 & 1 & 0
 \end{array}$$

- first effect on its column
- then look at  $c$
- finally look at  $b$

$$\begin{array}{c|cccccc}
 & x_1 & x_2 & x_3 & x_4 & -z & b \\
 \hline
 x_2 & 0 & (10 + \delta)1/5 + 4(-1/4) & 1/5 & -1/4 & 0 & 2 \\
 x_1 & 1 & (10 + \delta)(-1/5) + 4(1/2) & -1/5 & 1/2 & 0 & 8 \\
 \hline
 & 0 & -2/5\delta & -2/5 & -1 & 1 & -64
 \end{array}$$

- The dominant application of LP is mixed integer linear programming.
- In this context it is extremely important being able to begin with a model instantiated in one form followed by a sequence of problem modifications
  - row and column additions and deletions,
  - variable fixings

interspersed with resolves

- Derivation:
  1. economic interpretation
  2. bounding
  3. multipliers
  4. recipe
  5. Lagrangian
- Theory:
  - Symmetry
  - Weak duality theorem
  - Strong duality theorem
  - Complementary slackness theorem
- Dual Simplex
- Sensitivity Analysis, Economic interpretation