

DM559

Linear and Integer Programming

Lecture 8

Linear Transformations

Marco Chiarandini

Department of Mathematics & Computer Science
University of Southern Denmark

Outline

1. Linear Transformations
2. Coordinate Change
3. More on Change of Basis

Resume

- Linear dependence and independence
- Determine linear dependency of a set of vectors, ie, find non-trivial lin. combination that equal zero
- Basis
- Find a basis for a linear space
- Dimension (finite, infinite)

Outline

1. Linear Transformations
2. Coordinate Change
3. More on Change of Basis

Linear Transformations

Definition (Linear Transformation)

Let V and W be two vector spaces. A function $T : V \rightarrow W$ is **linear** if for all $\mathbf{u}, \mathbf{v} \in V$ and all $\alpha \in \mathbb{R}$:

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$
2. $T(\alpha\mathbf{u}) = \alpha T(\mathbf{u})$

A **linear transformation** is a linear function between two vector spaces

- If $V = W$ also known as **linear operator**
- Equivalent condition: $T(\alpha\mathbf{u} + \beta\mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v})$
- for all $\mathbf{0} \in V$, $T(\mathbf{0}) = \mathbf{0}$

Example (Linear Transformations)

- vector space $V = \mathbb{R}$, $F_1(x) = px$ for any $p \in \mathbb{R}$

$$\begin{aligned}\forall x, y \in \mathbb{R}, \alpha, \beta \in \mathbb{R} : F_1(\alpha x + \beta y) &= p(\alpha x + \beta y) = \alpha(px) + \beta(px) \\ &= \alpha F_1(x) + \beta F_1(y)\end{aligned}$$

- vector space $V = \mathbb{R}$, $F_2(x) = px + q$ for any $p, q \in \mathbb{R}$ or $F_3(x) = x^2$ are not linear transformations

$$T(x + y) \neq T(x) + T(y) \quad \text{for some } x, y \in \mathbb{R}$$

- vector spaces $V = \mathbb{R}^n$, $W = \mathbb{R}^m$, $m \times n$ matrix A , $T(\mathbf{x}) = A\mathbf{x}$ for $\mathbf{x} \in \mathbb{R}^n$

$$\begin{aligned}T(\mathbf{u} + \mathbf{v}) &= A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = T(\mathbf{u}) + T(\mathbf{v}) \\ T(\alpha\mathbf{u}) &= A(\alpha\mathbf{u}) = \alpha A\mathbf{u} = \alpha T(\mathbf{u})\end{aligned}$$

Example (Linear Transformations)

- vector spaces $V = \mathbb{R}^n$, $W : f : \mathbb{R} \rightarrow \mathbb{R}$. $T : \mathbb{R}^n \rightarrow W$:

$$T(\mathbf{u}) = T \left(\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} \right) = p_{u_1, u_2, \dots, u_n} = p_{\mathbf{u}}$$

$$p_{u_1, u_2, \dots, u_n} = u_1 x^1 + u_2 x^2 + u_3 x^3 + \dots + u_n x^n$$

$$p_{\mathbf{u}+\mathbf{v}}(x) = \dots = (p_{\mathbf{u}} + p_{\mathbf{v}})(x)$$

$$p_{\alpha \mathbf{u}}(\mathbf{x}) = \dots = \alpha p_{\mathbf{u}}(x)$$

Linear Transformations and Matrices

- any $m \times n$ matrix A defines a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m \rightsquigarrow T_A$
- for every linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ there is a matrix A such that $T(\mathbf{v}) = A\mathbf{v} \rightsquigarrow A_T$

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation and $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ denote the *standard basis* of \mathbb{R}^n and let A be the matrix whose columns are the vectors $T(\mathbf{e}_1), T(\mathbf{e}_2), \dots, T(\mathbf{e}_n)$: that is,

$$A = [T(\mathbf{e}_1) \ T(\mathbf{e}_2) \ \dots \ T(\mathbf{e}_n)]$$

Then, for every $\mathbf{x} \in \mathbb{R}^n$, $T(\mathbf{x}) = A\mathbf{x}$.

Proof: write any vector $\mathbf{x} \in \mathbb{R}^n$ as lin. comb. of standard basis and then make the image of it.

Example

$$T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

$$T \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} x + y + z \\ x - y \\ x + 2y - 3z \end{bmatrix}$$

- The image of $\mathbf{u} = [1, 2, 3]^T$ can be found by substitution:
 $T(\mathbf{u}) = [6, -1, -4]^T$.
- to find A_T :

$$T(\mathbf{e}_1) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad T(\mathbf{e}_2) = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \quad T(\mathbf{e}_3) = \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}$$

$$A = [T(\mathbf{e}_1) \ T(\mathbf{e}_2) \ T(\mathbf{e}_n)] = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 2 & -3 \end{bmatrix}$$

$$T(\mathbf{u}) = A\mathbf{u} = [6, -1, -4]^T.$$

Linear Transformation in \mathbb{R}^2


- We can visualize them!
- Reflection in the x axis:

$$T : \begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ -y \end{bmatrix} \quad A_T = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

- Stretching the plane away from the origin

$$T(\mathbf{x}) = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Rotation **anticlockwise** by an angle θ



we search the images of the standard basis vector $\mathbf{e}_1, \mathbf{e}_2$

$$T(\mathbf{e}_1) = \begin{bmatrix} a \\ c \end{bmatrix}, \quad T(\mathbf{e}_2) = \begin{bmatrix} b \\ d \end{bmatrix}$$

they will be orthogonal and with length 1.

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

For $\pi/4$:

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Identity and Zero Linear Transformations

- For $T : V \rightarrow V$ the linear transformation such that $T(\mathbf{v}) = \mathbf{v}$ is called the **identity**.
- if $V = \mathbb{R}^n$, the matrix $A_T = I$ (of size $n \times n$)
- For $T : V \rightarrow W$ the linear transformation such that $T(\mathbf{v}) = \mathbf{0}$ is called the **zero** transformation.
- If $V = \mathbb{R}^n$ and $W = \mathbb{R}^m$, the matrix A_T is an $m \times n$ matrix of zeros.

Composition of Linear Transformations

- Let $T : V \rightarrow W$ and $S : W \rightarrow U$ be linear transformations.
The **composition** of ST is again a linear transformation given by:

$$ST(\mathbf{v}) = S(T(\mathbf{v})) = S(\mathbf{w}) = \mathbf{u}$$

where $\mathbf{w} = T(\mathbf{v})$

- ST means do T and then do $S: V \xrightarrow{T} W \xrightarrow{S} U$
- if $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $S : \mathbb{R}^m \rightarrow \mathbb{R}^p$ in terms of matrices:

$$ST(\mathbf{v}) = S(T(\mathbf{v})) = S(A_T \mathbf{v}) = A_S A_T \mathbf{v}$$

note that composition is not commutative

Combinations of Linear Transformations

- If $S, T : V \rightarrow W$ are linear transformations between the same vector spaces, then $S + T$ and αS , $\alpha \in \mathbb{R}$ are linear transformations.
- hence also $\alpha S + \beta T$, $\alpha, \beta \in \mathbb{R}$ is

Inverse Linear Transformations

- If V and W are finite-dimensional vector spaces of the same dimension, then the **inverse** of a lin. transf. $T : V \rightarrow W$ is the lin. transf such that

$$T^{-1}(T(\mathbf{v})) = \mathbf{v}$$

- In \mathbb{R}^n if T^{-1} exists, then its matrix satisfies:

$$T^{-1}(T(\mathbf{v})) = A_{T^{-1}}A_T\mathbf{v} = I\mathbf{v}$$

that is, T^{-1} exists iff $(A_T)^{-1}$ exists and $A_{T^{-1}} = (A_T)^{-1}$
 (recall that if $BA = I$ then $B = A^{-1}$)

- In \mathbb{R}^2 for rotations:

$$A_{T^{-1}} = \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

Example

Is there an inverse to $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$

$$T \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} x + y + z \\ x - y \\ x + 2y - 3z \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 2 & -3 \end{bmatrix}$$

Since $\det(A) = 9$ then the matrix is invertible, and T^{-1} is given by the matrix:

$$A^{-1} = \frac{1}{9} \begin{bmatrix} 3 & 5 & 1 \\ 3 & -4 & 1 \\ 3 & -1 & -2 \end{bmatrix} \quad T^{-1} \left(\begin{bmatrix} u \\ v \\ w \end{bmatrix} \right) = \begin{bmatrix} \frac{1}{3}u + \frac{5}{9}v + \frac{1}{9}w \\ \frac{1}{3}u - \frac{4}{9}v + \frac{1}{9}w \\ \frac{1}{3}u + \frac{1}{9}v - \frac{2}{9}w \end{bmatrix}$$

Outline

1. Linear Transformations
2. Coordinate Change
3. More on Change of Basis

Coordinates

Recall:

Definition (Coordinates)

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis of a vector space V , then

- any vector $\mathbf{v} \in V$ can be expressed **uniquely** as $\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$
- and the real numbers $\alpha_1, \alpha_2, \dots, \alpha_n$ are the **coordinates** of \mathbf{v} wrt the basis S .

To denote the coordinate vector of \mathbf{v} in the basis S we use the notation

$$[\mathbf{v}]_S = \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{bmatrix}_S$$

- In the standard basis the coordinates of \mathbf{v} are precisely the components of the vector \mathbf{v} : $\mathbf{v} = v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + \dots + v_n \mathbf{e}_n$
- How to find coordinates of a vector \mathbf{v} wrt another basis?

Transition from Standard to Basis B

Definition (Transition Matrix)

Let $B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis of \mathbb{R}^n . The coordinates of a vector \mathbf{x} wrt B , $\mathbf{a} = [a_1, a_2, \dots, a_n]^T = [\mathbf{x}]_B$, are found by solving the linear system:

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n = \mathbf{x} \quad \text{that is} \quad \mathbf{x} = [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n][\mathbf{x}]_B$$

We call P the matrix whose columns are the basis vectors:

$$P = [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n]$$

Then for any vector $\mathbf{x} \in \mathbb{R}^n$

$$\mathbf{x} = P[\mathbf{x}]_B \quad \text{transition matrix from } B \text{ coords to standard coords}$$

moreover P is invertible (columns are a basis):

$$[\mathbf{x}]_B = P^{-1}\mathbf{x} \quad \text{transition matrix from standard coords to } B \text{ coords}$$

Example

$$B = \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \right\} \quad [\mathbf{v}]_B = \begin{bmatrix} 4 \\ 1 \\ -5 \end{bmatrix}$$

$$P = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 2 \\ -1 & 4 & 1 \end{bmatrix}$$

$\det(P) = 4 \neq 0$ so B is a basis of \mathbb{R}^3

We derive the standard coordinates of \mathbf{v} :

$$\mathbf{v} = 4 \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} - 5 \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -9 \\ -3 \\ -5 \end{bmatrix}$$

$$\mathbf{v} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 2 \\ -1 & 4 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 1 \\ -5 \end{bmatrix}_B = \begin{bmatrix} -9 \\ -3 \\ -5 \end{bmatrix}$$

Example (cntd)

$$B = \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \right\}, \quad [\mathbf{x}] = \begin{bmatrix} 5 \\ 7 \\ -3 \end{bmatrix}$$

We derive the B coordinates of vector \mathbf{x} :

$$\begin{bmatrix} 5 \\ 7 \\ -3 \end{bmatrix} = a_1 \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + a_2 \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} + a_3 \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

either we solve $P\mathbf{a} = \mathbf{x}$ in \mathbf{a} by Gaussian elimination or we find the inverse P^{-1} :

$$[\mathbf{x}]_B = P^{-1}\mathbf{x} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}_B \quad \text{check the calculation}$$

What are the B coordinates of the basis vector? ($[1, 0, 0]$, $[0, 1, 0]$, $[0, 0, 1]$)

Change of Basis

Since $T(\mathbf{x}) = P\mathbf{x}$ then $T(\mathbf{e}_i) = \mathbf{v}_i$, ie, T maps standard basis vector to new basis vectors

Example

Rotate basis in \mathbb{R}^2 by $\pi/4$ anticlockwise, find coordinates of a vector wrt the new basis.

$$A_T = \begin{bmatrix} \cos \frac{\pi}{4} & -\sin \frac{\pi}{4} \\ \sin \frac{\pi}{4} & \cos \frac{\pi}{4} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Since the matrix A_T rotates $\{\mathbf{e}_1, \mathbf{e}_2\}$, then $A_T = P$ and its columns tell us the coordinates of the new basis and $\mathbf{v} = P[\mathbf{v}]_B$ and $[\mathbf{v}]_B = P^{-1}\mathbf{v}$. The inverse is a rotation **clockwise**:

$$P^{-1} = \begin{bmatrix} \cos(-\frac{\pi}{4}) & -\sin(-\frac{\pi}{4}) \\ \sin(-\frac{\pi}{4}) & \cos(-\frac{\pi}{4}) \end{bmatrix} = \begin{bmatrix} \cos(\frac{\pi}{4}) & \sin(\frac{\pi}{4}) \\ -\sin(\frac{\pi}{4}) & \cos(\frac{\pi}{4}) \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Example (cntd)

Find the new coordinates of a vector $\mathbf{x} = [1, 1]^T$

$$[\mathbf{x}]_B = P^{-1}\mathbf{x} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix}$$

Change of basis from B to B'

Given a basis B of \mathbb{R}^n with transition matrix P_B ,
 and another basis B' with transition matrix $P_{B'}$,
 how do we change from coords in the basis B to coords in the basis B' ?

coordinates in $B \xrightarrow{v=P_B[v]_B}$ standard coordinates $\xrightarrow{[v]_{B'}=P_{B'}^{-1}v}$ coordinates in B'

$$[v]_{B'} = P_{B'}^{-1} P_B [v]_B$$

$$M = P_{B'}^{-1} P_B = P_{B'}^{-1} [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n] \stackrel{\text{ex11sh2}}{=} [P_{B'}^{-1} \mathbf{v}_1 \ P_{B'}^{-1} \mathbf{v}_2 \ \dots \ P_{B'}^{-1} \mathbf{v}_n]$$

Theorem

If B and B' are two bases of \mathbb{R}^n , with

$$B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$$

then the transition matrix from B coordinates to B' coordinates is given by

$$M = [[\mathbf{v}_1]_{B'} \ [\mathbf{v}_2]_{B'} \ \dots \ [\mathbf{v}_n]_{B'}]$$

(the columns of M are the B' coordinates of the basis B)

Example

$$B = \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} \quad B' = \left\{ \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 5 \\ 2 \end{bmatrix} \right\}$$

are basis of \mathbb{R}^2 , indeed the corresponding transition matrices from standard basis:

$$P = \begin{bmatrix} 1 & -1 \\ 2 & 1 \end{bmatrix} \quad Q = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$$

have $\det(P) = 3$, $\det(Q) = 1$. Hence, lin. indep. vectors.

We are given

$$[\mathbf{x}]_B = \begin{bmatrix} 4 \\ -1 \end{bmatrix}_B$$

find its coordinates in B' .

Example (cntd)

1. find first the standard coordinates of \mathbf{x}

$$\mathbf{x} = 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ -1 \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

and then find B' coordinates:

$$[\mathbf{x}]_S = Q^{-1}\mathbf{x} = \begin{bmatrix} 2 & -5 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 5 \\ 7 \end{bmatrix} = \begin{bmatrix} -25 \\ 16 \end{bmatrix}_S$$

2. use transition matrix M from B to B' coordinates:

$$\mathbf{v} = P[\mathbf{v}]_B \quad \text{and} \quad \mathbf{v} = Q[\mathbf{v}]_{B'} \quad \rightsquigarrow \quad [\mathbf{v}]_{B'} = Q^{-1}P[\mathbf{v}]_B:$$

$$M = Q^{-1}P = \begin{bmatrix} 2 & -5 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} -8 & -7 \\ 5 & 4 \end{bmatrix}$$

$$[\mathbf{x}]_{B'} = \begin{bmatrix} -8 & -7 \\ 5 & 4 \end{bmatrix} \begin{bmatrix} 4 \\ -1 \end{bmatrix} = \begin{bmatrix} -25 \\ 16 \end{bmatrix}_{B'}$$

Outline

1. Linear Transformations
2. Coordinate Change
3. More on Change of Basis

Change of Basis for a Lin. Transf.

We saw how to find A for a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ using standard basis in both \mathbb{R}^n and \mathbb{R}^m . Now: is there a matrix that represents T wrt two arbitrary bases B and B' ?

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation and $B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ and $B' = \{\mathbf{v}'_1, \mathbf{v}'_2, \dots, \mathbf{v}'_m\}$ be bases of \mathbb{R}^n and \mathbb{R}^m . Then for all $\mathbf{x} \in \mathbb{R}^n$, $[T(\mathbf{x})]_{B'} = M[\mathbf{x}]_B$ where $M = A_{[B, B']}$ is the $m \times n$ matrix with the i th column equal to $[T(\mathbf{v}_i)]_{B'}$, the coordinate vector of $T(\mathbf{v}_i)$ wrt the basis B' .

Proof:

change B to standard	$\mathbf{x} = P_B^{n \times n}[\mathbf{x}]_B \quad \forall \mathbf{x} \in \mathbb{R}^n$
↓	
perform linear transformation in standard coordinates	$T(\mathbf{x}) = A\mathbf{x} = AP_B^{n \times n}[\mathbf{x}]_B$
↓	
change to basis B'	$[\mathbf{u}]_{B'} = (P_{B'}^{m \times m})^{-1}\mathbf{u} \quad \forall \mathbf{u} \in \mathbb{R}^m$ $[T(\mathbf{x})]_{B'} = (P_{B'}^{m \times m})^{-1}AP_B^{n \times n}[\mathbf{x}]_B$ $M = (P_{B'}^{m \times m})^{-1}AP_B^{n \times n}$

How is M done?

- $P_B = [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n]$
- $AP_B = A[\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n] = [A\mathbf{v}_1 \ A\mathbf{v}_2 \ \dots \ A\mathbf{v}_n]$
- $A\mathbf{v}_i = T(\mathbf{v}_i)$: $AP_B = [T(\mathbf{v}_1) \ T(\mathbf{v}_2) \ \dots \ T(\mathbf{v}_n)]$
- $M = P_{B'}^{-1}AP_B = [P_{B'}^{-1}T(\mathbf{v}_1) \ P_{B'}^{-1}T(\mathbf{v}_2) \ \dots \ P_{B'}^{-1}T(\mathbf{v}_n)]$
- $M = [[T(\mathbf{v}_1)]_{B'} \ [T(\mathbf{v}_2)]_{B'} \ \dots \ [T(\mathbf{v}_n)]_{B'}]$

Hence, if we change the basis from the standard basis of \mathbb{R}^n and \mathbb{R}^m the matrix representation of T changes

Similarity

Particular case $m = n$:

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation
and $B = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ be a basis \mathbb{R}^n .

Let A be the matrix corresponding to T in standard coordinates: $T(\mathbf{x}) = A\mathbf{x}$.

Let

$$P = [\mathbf{x}_1 \ \mathbf{x}_2 \ \cdots \ \mathbf{x}_n]$$

be the matrix whose columns are the vectors of B . Then for all $\mathbf{x} \in \mathbb{R}^n$,

$$[T(\mathbf{x})]_B = P^{-1}AP[\mathbf{x}]_B$$

Or, the matrix $A_{[B,B]} = P^{-1}AP$ performs the same linear transformation as the matrix A but expressed it in terms of the basis B .

Similarity

Definition



A square matrix C is **similar** (represent the same linear transformation) to the matrix A if there is an invertible matrix P such that

$$C = P^{-1}AP.$$

Similarity defines an equivalence relation:

- (reflexive) a matrix A is similar to itself
- (symmetric) if C is similar to A , then A is similar to C
 $C = P^{-1}AP$, $A = Q^{-1}CQ$, $Q = P^{-1}$
- (transitive) if D is similar to C , and C to A , then D is similar to A

Example



- $x^2 + y^2 = 1$ circle in standard form
- $x^2 + 4y^2 = 4$ ellipse in standard form
- $5x^2 + 5y^2 - 6xy = 2$??? Try rotating $\pi/4$ anticlockwise

$$A_T = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = P$$

$$\mathbf{v} = P[\mathbf{v}]_B \iff \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix}$$

$$X^2 + 4Y^2 = 1$$

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$:

$$T \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x + 3y \\ -x + 5y \end{bmatrix}$$

What is its effect on the xy -plane?

Let's change the basis to

$$B = \{\mathbf{v}_1, \mathbf{v}_2\} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}$$

Find the matrix of T in this basis:

- $C = P^{-1}AP$, A matrix of T in standard basis, P is transition matrix from B to standard

$$C = P^{-1}AP = \frac{1}{2} \begin{bmatrix} -1 & 3 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 2 \end{bmatrix}$$

Example (cntd)

- the B coordinates of the B basis vectors are

$$[\mathbf{v}_1]_B = \begin{bmatrix} 1 \\ 0 \end{bmatrix}_B, \quad [\mathbf{v}_2]_B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}_B$$

- so in B coordinates T is a stretch in the direction \mathbf{v}_1 by 4 and in dir. \mathbf{v}_2 by 2:

$$[T(\mathbf{v}_1)]_B = \begin{bmatrix} 4 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}_B = \begin{bmatrix} 4 \\ 0 \end{bmatrix}_B = 4[\mathbf{v}_1]_B$$

- The effect of T is however the same no matter what basis, only the matrices change! So also in the standard coordinates we must have:

$$A\mathbf{v}_1 = 4\mathbf{v}_1 \quad A\mathbf{v}_2 = 2\mathbf{v}_2$$

Summary

- Linear transformations and proofs that a given mapping is linear
- two-way relationship between matrices and linear transformations
- change from standard (S) to arbitrary basis (B)
- change of basis between two arbitrary basis (from B to B')
- Matrix representation of a transformation with respect to two arbitrary basis
- Similarity of square matrices