

DM545/DM871 – Linear and integer programming

Sheet 2, Spring 2019 [pdf format]

Exercise 1 List all vertices of the polyhedron $Ax \leq b$ characterized by the following matrices A and b :

$$A = \begin{bmatrix} 2 & 0 & 1 & -4 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 0 \end{bmatrix} \quad b = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$$

You are encouraged to use Python for carrying out the calculations.

Exercise 2* Simplex method

This is part of the first exercise (Opgave 1) in the written exam of 2008. Consider the following linear programming problem (P1)

$$\begin{aligned} & \text{maximize} && 2x_1 + 4x_2 - x_3 \\ & \text{subject to} && 2x_1 - x_3 \leq 6 \\ & && 3x_2 - x_3 \leq 9 \\ & && x_1 + x_2 \leq 4 \\ & && x_1, x_2, x_3 \geq 0 \end{aligned}$$

- Rewrite the problem in equational standard form adding the slack variables x_4, x_5, x_6 to the three constraints above, respectively, and write the first simplex tableau with x_4, x_5, x_6 as basic solution.
- Argue that x_2 can be brought in the basis with advantage and perform one pivot iteration that brings x_2 into the basic solution.
- After another pivot iteration, it is x_1 that can be brought with advantage in the basis (you do not have to perform this iteration), reaching the following simplex tableau:

x1	x2	x3	x4	x5	x6	-z	b
0	0	-5/3	1	2/3	-2	0	4
0	1	-1/3	0	1/3	0	0	3
1	0	1/3	0	-1/3	1	0	1
0	0	-1/3	0	-2/3	-2	1	-14

Argue that an optimal solution is found and give the solution together with its objective value.

Exercise 3* Simplex method

Solve the following LP problem carrying out the simplex operations:

$$\begin{aligned} & \text{maximize} && 5x_1 + 4x_2 + 3x_3 \\ & \text{subject to} && 2x_1 + 3x_2 + x_3 \leq 5 \\ & && 4x_1 + x_2 + 2x_3 \leq 11 \\ & && 3x_1 + 4x_2 + 2x_3 \leq 8 \\ & && x_1, x_2, x_3 \geq 0 \end{aligned}$$

You are free to use any of the two representations, tableau or dictionary.

You can also get help from Python. You find a tutorial for what you need at this link:




Figure 1: A network with activities on nodes for a small project with 6 activities. For each activity the following data is given in that order from left to right: normal time, minimum time in weeks, and the cost of shortening the duration of the activity by one week.

<http://www.imada.sdu.dk/~marco/DM545/Resources/Ipython/Tutorial4Exam.html>.

Exercise 4

Solve the following linear programming problem applying the simplex algorithm:

$$\begin{aligned}
 &\text{maximize} && 3x_1 + 2x_2 \\
 &\text{subject to} && x_1 - 2x_2 \leq 1 \\
 & && x_1 - x_2 \leq 2 \\
 & && 2x_1 - x_2 \leq 6 \\
 & && x_1 \leq 5 \\
 & && 2x_1 + x_2 \leq 16 \\
 & && x_1 + x_2 \leq 12 \\
 & && x_1 + 2x_2 \leq 21 \\
 & && x_2 \leq 10 \\
 & && x_1, x_2 \geq 0.
 \end{aligned}$$

[Hint: you can plot the feasibility region with one of the tools linked at the course web page: “Tools” -> “Web applications on the simplex” -> “LP Simplex” and use the clairvoyant’s rule to minimize the number of operations to carry out.]

Exercise 5* Project Scheduling

This exercise is a part of one that appeared in Exam 2011.

A small project has 6 sub-activities A, B, C, D, E, F whose individual dependency (shown by the immediate predecessors) is given in Figure 1. Here we also list the normal time (in weeks), the absolute minimum time and the cost of shortening the activity by one week.

The goal is to shorten the duration of the project to 19 weeks. This means that the duration of one or more activities has to be shortened. Of course we want to select these so that the total cost of shortening the duration to 19 weeks is minimized. Formulate this problem as a linear programming problem and argue that the optimal solution to this LP will provide the correct answer. Note that you must use the actual data in the LP formulation!

Exercise 6 Quizzes

1. In 4D, how many hyperplanes need to intersect to give a point?
2. In 4D, can a point be described by more than 4 hyperplanes?
3. Consider the intersection of n hyperplanes in n dimensions: when does it uniquely identify a point?

Vertices of Polyhedra:

Consider the polyhedron described by $Ax \leq \mathbf{b}$, $A \in \mathbb{R}^{m \times n}$, $\mathbf{x} \in \mathbb{R}^n$, that is:

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &\leq b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &\leq b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &\leq b_m \end{aligned}$$

4. For a point \mathbf{x} of a polyhedron, we define as *active* constraints those that are satisfied to equality by \mathbf{x} . How many constraints are *active* in a *vertex* of a polyhedron $Ax \leq \mathbf{b}$, $A \in \mathbb{R}^{m \times n}$, $\mathbf{x} \in \mathbb{R}^n$?
5. Does every point \mathbf{x} that activates n constraints form a vertex of the polyhedron?
6. Can a vertex activate more than n constraints?
7. What if there are more variables than constraints? If $m > n$ then we can find a subset and then activate but what if $m < n$, can we have a vertex?
8. Combinatorial explosion of vertices: how many constraints and vertices has an n -dimensional hypercube?
9. If there are m constraints and n variables, $m > n$, what is an upper bound to the number of vertices?

Tableaux and Vertices

10. For each of these three statements, say if they are true or false:

- One tableau \implies one vertex of the feasible region
- One tableau \longleftarrow one vertex of the feasible region
- One tableau \iff one vertex of the feasible region

11. Consider the following LP problem and the corresponding final tableau:

$$\begin{array}{ll} \max & 6x_1 + 8x_2 \\ & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0 \end{array} \quad \begin{array}{c|cccc|c} & x_1 & x_2 & x_3 & x_4 & -z & b \\ \hline x_2 & 0 & 1 & 1/5 & -1/4 & 0 & 2 \\ x_1 & 1 & 0 & -1/5 & 1/2 & 0 & 8 \\ \hline & 0 & 0 & -2/5 & -1 & 1 & -64 \end{array}$$

- How many variables (original and slack) can be different from zero?
- $(x_3, x_4) = (0, 0)$ are non basic, what does this tell us about the constraints?

Let's generalize the previous case. Consider an LP with m constraints, n original variables and m slack variables. In an optimal solution:

- if $m > n$, how many variables (original and slack) can be nonzero at most?
- if $m < n$ how many original variables must be zero at least? In other terms, in a mix planning problem with n products and m , $m < n$ resources, how many products at most will be to be produced in an optimal solution?

12. Consider the following LP problem and the corresponding final tableau:

$$\begin{array}{ll} \max & 6x_1 + 8x_2 \\ & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0 \end{array} \quad \begin{array}{c|cccc|c} & x_1 & x_2 & x_3 & x_4 & -z & b \\ \hline x_3 & 0 & 0 & 1 & 1/2 & 0 & 1 \\ x_1 & 1 & 1 & 0 & -1/2 & 0 & 1 \\ \hline & 0 & -2 & 0 & 1/2 & 1 & -1 \end{array}$$

$(x_2, x_4) = (0, 0)$ is non basic, what does this tell us about the constraints?

13. If in the original space of the problem we had 3 variables, and there are 6 constraints, how many constraints would be active?
14. For the general case with n original variables:
One basic feasible solution \iff a matrix of active constraints has rank n . True or False?
15. Consider an LP problem with m constraints and n original variables, $m > n$. We saw that in \mathbb{R}^n a point is the intersection of at least n hyperplanes. In LP this corresponds to say that in a vertex there are n active constraints. Let a tableau be associated with a solution that makes exactly $n + 1$ constraints active, what can we say about the corresponding basic and non-basic variable values?
16. Given a polyhedron, what is the algebraic definition of vertex adjacency in 2, 3 and n dimensions?
How does this condition translate in terms of tableau?

Exercise 7


What argument is used to prove that the simplex algorithm always terminates in a finite number of iterations if it does not encounter a situation in which one of the basic variables is zero? What may happen instead if the latter situation arises and which remedies are introduced?

Exercise 8*

Exercise 3 from Exam 2013.

Consider the following LP problem:

$$(P) \quad \begin{array}{ll} \max & z = x_1 + 5x_2 \\ \text{s.t.} & -x_1 + 3x_2 \leq 6 \\ & 4x_1 + 4x_2 \geq 5 \\ & 0 \leq x_1 \leq 2 \\ & x_2 \geq 0 \end{array}$$



(Note: the following subtasks can be carried out independently; use fractional mode for numerical calculations; in the online version you find the problem in ASCII format).

- a. The polyhedron representing the feasibility region is depicted in the figure. Indicate for each of the four points represented whether they are feasible and/or basic solutions. Justify your answer.
- b. Write the initial tableau or dictionary for the simplex method. Write the corresponding basic solution and its value. State whether the solution is feasible or not and whether it is optimal or not.
- c. Consider the following tableau:

	x1	x2	x3	x4	x5	-z	b
I	0	4	1	-1/4	0	0	29/4
II	1	1	0	-1/4	0	0	5/4
III	0	-1	0	1/4	1	0	3/4
IV	0	4	0	1/4	0	1	-5/4

and the following three pivoting rules:

- largest coefficient
- largest increase
- steepest edge.

Which entering and leaving variables would each of them indicate? In this specific case, which rule would be convenient to follow? Report the details of the computations for the first two rules and carry out graphically the application of the third rule using the plot in the figure above (tikz code to reproduce the figure available in the online version.)

Exercise 9*

The two following LP problems lead to two particular cases when solved by the simplex algorithm. Identify these cases and characterize them, that is, give indication of which conditions generate them in general.

$$\begin{array}{ll} \text{maximize} & 2x_1 + x_2 \\ \text{subject to} & x_2 \leq 5 \\ & -x_1 + x_2 \leq 1 \\ & x_1, x_2 \geq 0 \end{array}$$

$$\begin{array}{ll} \text{maximize} & x_1 + x_2 \\ \text{subject to} & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0 \end{array}$$

Exercise 10*

Consider the following problem:

$$\begin{array}{ll} \max & z = 4x_2 \\ \text{s.t.} & 2x_2 \geq 0 \\ & -3x_1 + 4x_2 \geq 1 \\ & x_1, x_2 \geq 0 \end{array}$$

- a. Write the LP in equational standard form and say why it does not provide immediately an initial feasible basis for the simplex method.
- b. To overcome the situation of infeasible basis construct the auxiliary problem for a phase I–phase II solution approach. Determine which variables are initially in basis and which are not in basis in the auxiliary problem.
- c. Answer the following questions
 - i) Is the initial basis in the auxiliary problem feasible in the original problem?
 - ii) Is it optimal in the auxiliary problem?
 - iii) Is it degenerate?
 - iv) Can we say at this stage if phase I will terminate?
 - v) If it will terminate, can we say at this stage that it will terminate with a basis that corresponds to a feasible solution in the original problem?
 - vi) Solve the problem by carrying out Phase I and Phase II of the simplex algorithm.