

DM534 — Øvelser Uge 46

Introduktion til Datalogi, Efterår 2021

Jonas Vistrup

1 I

1.1

Which of the following formulas are satisfiable (give a satisfying assignment)? Which are not (give reasons)?

- (a) $A \wedge B$. **SVAR:** Satisfiable, $A=t$ $B=t$.
- (b) $A \vee B$. **SVAR:** Satisfiable, $A=t$ $B=t$.
- (c) $A \Rightarrow B$. **SVAR:** Satisfiable, $A=t$ $B=t$.
- (d) $A \wedge \neg A$. **SVAR:** Unsatisfiable.
- (e) $A \vee \neg A$. **SVAR:** Satisfiable, $A=t$.

1.2

Two formulas are equivalent, if the same assignments satisfy both of them.

Which of the following formulas are equivalent?

- (a) $\neg A \wedge B$.
- (b) $\neg A \vee B$.
- (c) $A \Rightarrow B$.
- (d) $(A \Rightarrow B) \wedge (\neg B \Rightarrow A)$.
- (e) $(\neg A \Rightarrow B) \wedge (\neg B \Rightarrow \neg A)$.

SVAR: (b) \Leftrightarrow (c) og (d) \Leftrightarrow (e).

1.3

Convert the following formulas into CNF:

- (a) $\neg A \wedge B$. **SVAR:** $(\neg A) \wedge (B)$.
- (b) $\neg A \vee B$. **SVAR:** $(\neg A \vee B)$.
- (c) $A \Rightarrow B$. **SVAR:** $(\neg A \vee B)$.

(d) $(A \Rightarrow B) \wedge (-B \Rightarrow A)$. **SVAR:** $(-A \vee B) \wedge (B \vee A)$.

1.4

- (a) Write two clauses that forbid solutions where there is a queen in the right half of the first row. **SVAR:** $(\neg X_{1,1}) \wedge (\neg X_{1,2})$
- (b) Instead of adding two clauses, change an existing clause. **SVAR:** Change $(X_{1,1} \vee X_{1,2} \vee X_{1,3} \vee X_{1,4})$ to $(X_{1,3} \vee X_{1,4})$.

1.5

Run SAT solver.

1.6

The formula from Slide 11 contains redundant information. For example, $X_{1,1} \Rightarrow -X_{1,2}$ and $X_{1,2} \Rightarrow -X_{1,1}$ are equivalent. Understand and remove these redundancies:

- (a) Why do these redundancies occur?
- (b) Identify all such redundancies!
- (c) Write down a simplified formula without redundancies!
- (d) Convert the simplified formula into CNF!
- (e) Write the formula in DIMACS format!
- (f) Run the lingeling solver on it and interpret the result!

SVAR a: A tower in spot x might exclude a tower in spot y, and the same is true reverse, but both of these statements are equivalent to stating, that a tower can't be in both x and y.

SVAR b:

- $X_{1,1} \Rightarrow -X_{1,2}$ og $X_{1,2} \Rightarrow -X_{1,1}$.
- $X_{1,1} \Rightarrow -X_{2,1}$ og $X_{2,1} \Rightarrow -X_{1,1}$.
- $X_{1,2} \Rightarrow -X_{2,2}$ og $X_{2,2} \Rightarrow -X_{1,2}$.
- $X_{2,1} \Rightarrow -X_{2,2}$ og $X_{2,2} \Rightarrow -X_{2,1}$.

SVAR c:

$$(X_{1,1} \Rightarrow -X_{1,2}) \wedge (X_{1,1} \Rightarrow -X_{2,1}) \wedge (X_{1,2} \Rightarrow -X_{2,2}) \wedge (X_{2,1} \Rightarrow -X_{2,2}) \\ \wedge (X_{1,1} \vee X_{1,2}) \wedge (X_{2,1} \vee X_{2,2})$$

SVAR d:

$$(-X_{1,1} \vee -X_{1,2}) \wedge (-X_{1,1} \vee -X_{2,1}) \wedge (-X_{1,2} \vee -X_{2,2}) \wedge (-X_{2,1} \vee -X_{2,2})$$

$$\wedge(X_{1,1} \vee X_{1,2}) \wedge (X_{2,1} \vee X_{2,2})$$

SVAR e: p cnf 4 6 -1 -2 0 -1 -3 0 -2 -4 0 -3 -4 0 1 2 0 3 4 0

SVAR f: Did in 1.5. Got 1, -2, -3, 4. This translates to a tower on $X_{1,1}$ and $X_{2,2}$.

2 II

2.1

Which of the following formulas are satisfiable (give a satisfying assignment)? Which are not (give reasons)?

- (a) $(A \Rightarrow B) \wedge (B \Rightarrow A)$. **SVAR:** Satisfiable, A=t B=t.
- (b) $(A \Rightarrow B) \wedge (B \Rightarrow A) \wedge A$. **SVAR:** Satisfiable, A=t B=t.
- (c) $(A \Rightarrow B) \wedge (B \Rightarrow A) \wedge \neg A$. **SVAR:** Satisfiable, A=f B=f.
- (d) $(A \Rightarrow B) \wedge (B \Rightarrow A) \wedge (\neg A \Rightarrow \neg B) \wedge (\neg B \Rightarrow A)$. **SVAR:** Satisfiable, A=t B=t.

2.2

Two formulas are equivalent, if the same assignments satisfy both of them.

Which of the following formulas are equivalent?

- (a) $(A \Rightarrow B) \wedge (\neg B \Rightarrow A)$.
- (b) $(A \Rightarrow \neg B) \wedge (B \Rightarrow A)$.
- (c) $(\neg A \vee \neg B) \wedge (A \vee \neg B)$.
- (d) $(B \vee A) \wedge (\neg A \vee B)$.

SVAR: (a) \Leftrightarrow (d) og (b) \Leftrightarrow (c).

2.3

Convert the following formulas into CNF:

- (a) $(\neg A \Rightarrow B) \wedge (\neg B \Rightarrow \neg A)$. **SVAR:** $(A \vee B) \wedge (B \vee \neg A)$.
- (b) $A \Rightarrow (\neg(B \wedge D))$. **SVAR:** $(\neg A \vee \neg B \vee \neg D)$
- (c) $A \Rightarrow (\neg(B \vee D))$. **SVAR:** $(\neg A \vee \neg B) \wedge (\neg A \vee \neg D)$
- (d) $A \Rightarrow (\neg(B \Rightarrow (C \wedge D)))$. **SVAR:** $(\neg A \vee \neg B) \wedge (\neg A \vee \neg C \vee \neg D)$.

2.4

Write a Java program that generates the input for a SAT solver to solve the 3-Towers problem:

- (a) Write a method `pair2int(int r, int c)`. Should map (1,1), (1,2), ..., (3,3) to 1 to 9 using the formula $3 \cdot (r - 1) + c$.
- (b) Write nested for-loops that go through all positions on the board from (1,1) to (3,3) and produces clauses that represent attacks.

- (c) Write a for-loop that produces clauses that specify that all 3 rows contain a tower.
- (d) Using (a)–(c), write a DIMACS file and test it using `lingeling`.

2.5

Generalize your Java program from Exercise II-4 to generate the input for a SAT solver to solve the N-Towers problem:

- (a) The method `pair2int(int n,int r,int c)` should map pairs (r,c) to the integers 1 to n^2 using the formula $n \cdot (r - 1) + c$.
- (b) Write nested for-loops that go through all positions on the board from $(1,1)$ to (n,n) and produces clauses that represent attacks.
- (c) Write a for-loop that produces clauses that specify that all rows contain a tower.
- (d) Using (a)–(c), write a DIMACS file and test it using `lingeling`.

2.6

Extend your Java program from Exercise II-5 to generate the input for a SAT solver to solve the N-Queens problem:

- (a) Reuse your function `pair2int(n,r,c)` from Exercise II-5.
- (b) Adapt your for-loops from Exercise II-5 to produce also clauses for the diagonals.
- (c) Reuse the for-loop from Exercise II-5 that produces clauses that specify that all rows contain a tower.
- (d) Using (a)–(c), write a DIMACS file and test it using `lingeling`.