

# DM534 — Øvelser Uge ??

Introduktion til Datalogi, Efterår 2021


Jonas Vistrup, med tilføjelser af Rolf Fagerberg

---


## 1 I

### 1.1

Draw the graph representing the road system in the figure below, and write down the number of vertices, the number of edges and the degree of each vertex.



SVAR:




5 vertices and 7 edges.

## 1.2

On Twitter:

- John follows Joan, Jean and Jane; Joe follows Jane and Joan; Jean and Joan follow each other. Draw a digraph illustrating these follow-relationships between John, Joan, Jean, Jane and Joe.
- Twitter has  $\approx 313$  million active users (June 2016, based on Twitter Inc.). Imagine you would like to store the digraph for the follow-relationships in an adjacency matrix that uses 4 bytes per entry on your new laptop which has 64 GB of RAM. Is this feasible?
- The municipality of Odense has a population of  $\approx 200000$  people. Let  $G$  be the graph where the meaning of an edge from vertex  $i$  to  $j$  is "person  $i$  is friends with person  $j$ ". Imagine you would like to store the adjacency matrix for this graph for the relationships in a matrix representation that uses 4 bytes per entry on your new laptop which has 64 GB of RAM. Is this feasible?

**SVAR a:**




**SVAR b:**  $(313 \cdot 10^6)^2 \cdot 4 \text{ bytes} = 3.91876 \cdot 10^{17} \text{ bytes} > 64 \cdot 10^9 \text{ bytes}$ . No.

**SVAR c:**  $(2 \cdot 10^5)^2 \cdot 4 \text{ bytes} = 160 \cdot 10^9 \text{ bytes} > 64 \cdot 10^9 \text{ bytes}$ . No.

## 1.3

Consider the following six graphs (note that the nodes do not have labels).



- (a) How many walks of length 3 from the red vertex to the green vertex are there in graph 3?
- (b) How many paths from the red vertex to the green vertex are there in graph 3?
- (c) How many shortest paths from the red vertex to the green vertex are there in graph 3?
- (d) For each of the graphs: what is the longest of all pairwise shortestpaths?
- (e) Give an adjacency matrix for graph 1. Can there be different adjacency matrices for the same graph? If so, name a second adjacency matrix for graph 1. Can you find two different adjacency matrices for graph 6? All orders of vertex resulting in the same adjacency matrix for graph 6.

**SVAR a:** 4.

**SVAR b:** 2.

**SVAR c:** 1.

**SVAR d:** 2,3,2,2,1.

**SVAR e:**

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$$

## 1.4

Let  $A$  be an adjacency matrix. In the lecture you learned that the  $ij$ -entry of  $A^k$  is the number of different walks from vertex  $i$  to vertex  $j$  using exactly  $k$  edges.


- (a) What is the interpretation of  $ij$ -entry of the matrix  $A^1 + A^2 + A^3$ ?
- (b) Complete the following sentence with the missing expression: In a graph  $G$  with adjacency matrix  $A$ , vertex  $i$  and  $j$  ( $i \neq j$ ) are connected if and only if ...  $> 0$ .

**SVAR a:** The number of walks from  $i$  to  $j$  with length 1, 2 or 3.

**SVAR b:** They are connected if and only if  $c_{ij} > 0$ , where  $c_{ij}$  is the entry in the  $i$ 'th row and  $j$ 'th column in the matrix  $C = A^1 + A^2 + \dots + A^{n-1}$ . It is enough to consider powers up to  $k - 1$ , since if  $i$  og  $j$  are connected by a walk, they are also connected by a path, i.e., a walk without repetitions among the nodes (a repetition will mean that the walk contains a cycle, which can just be removed from the walk, after which it will still connect  $i$  and  $j$ ). If we want the statement to also hold for  $i = j$ , we can just add  $A^0$  to the sum ( $A^0$  is defined to be the identity matrix, i.e., the matrix having ones on the diagonal and zeros elsewhere).

## 1.5

Let the following weighted graph (from the lecture slides, weights are depicted in red) be given:



It has the following distance matrix  $D$ :

$$D = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{matrix} & \begin{pmatrix} 0 & 1 & 3 & 7 & 2 & 6 \\ 1 & 0 & 2 & 6 & 3 & 5 \\ 3 & 2 & 0 & 4 & 5 & 3 \\ 7 & 6 & 4 & 0 & 6 & 1 \\ 2 & 3 & 5 & 6 & 0 & 7 \\ 6 & 5 & 3 & 1 & 7 & 0 \end{pmatrix} \end{matrix}$$

- How many shortest path in  $G$  are of length 6? Name them. **SVAR:** 1-6, 2-4 and 4-5.
- How long is the longest of all pairwise shortest paths in the graph? Are there several longest shortest paths? **SVAR:** 1-4 and 5-6 are both 7 long.
- How many paths in  $G$  are of length 6? (Note: a path does not necessarily need to be a shortest path.) Name them. **SVAR:** 1-2, 1-6, 2-4, 3-5, 4-5.


## 1.6

Assume in this exercise that all weights on edges are non-negative values.


- In a graph  $G$  with  $n = 6$  vertices, how many matrix-matrix multiplication operations are needed in the worst case in order to compute the distance matrix  $D$ , when the method of repeated squaring is used to compute  $D$ ? **SVAR:**  $\lceil \log_2(6) \rceil = 3$
- In a graph  $G$  with  $n = 200$  vertices, how many matrix-matrix multiplications are needed in the worst case in order to compute the distance matrix  $D$ , when the method of repeated squaring is used to compute  $D$ ? **SVAR:**  $\lceil \log_2(200) \rceil = 8$

- (c) Can you find a graph  $G$  with  $n = 6$  vertices, for which  $W^4 \neq W^5$ ? If so, depict it.
- (d) Can you find a graph  $G$  with  $n = 6$  vertices, for which  $W^5 \neq W^6$ ? If so, depict it.  
**SVAR:** No, after each step at least one new vertex is visited if there is a change.
- (e) Can you find a graph  $G$  with  $n = 6$  vertices, for which  $W^1 = W^2$ ? If so, depict it.
- (f) What is the computational runtime in order to compute the distance matrix  $D$  for a graph  $G$  with  $n$  vertices if the method of repeated squaring is used to compute  $D$ ?  
**SVAR:**  $n^3 \log_2(n)$ .

**SVAR c:**




**SVAR e:**



## 2 II

### 2.1


Consider the following molecule (it's called 2,3-Dimethylhexane, see <https://en.wikipedia.org/wiki/2,3-Dimethylhexane>):



- How many carbon atoms does this molecule have?
- Draw the graph  $G$  corresponding to the carbon backbone of the molecule.
- Give the edge weight matrix  $W$  for the graph  $G$ .
- Use your brain or the Java program ShortestPaths.java to infer the distance matrix (Hint: the graph is rather simple, you won't need a program for that.)
- What is the Wiener Index  $\mathcal{W}(G)$ ?
- How many shortest paths of length 3  $i \rightarrow \dots \rightarrow j$  with  $i < j$  are in  $G$ ?
- Using Wiener's method for predicting the boiling point, what is your prediction for 2,3-Dimethylhexane?

**SVAR a:** 8.

**SVAR b:**



**SVAR c:**

$$\begin{pmatrix} 0 & 1 & \infty & \infty & \infty & \infty & \infty & \infty \\ 1 & 0 & 1 & 1 & \infty & \infty & \infty & \infty \\ \infty & 1 & 0 & \infty & \infty & \infty & \infty & \infty \\ \infty & 1 & \infty & 0 & 1 & 1 & \infty & \infty \\ \infty & \infty & \infty & 1 & 0 & \infty & \infty & \infty \\ \infty & \infty & \infty & 1 & \infty & 0 & 1 & \infty \\ \infty & \infty & \infty & \infty & \infty & 1 & 0 & 1 \\ \infty & \infty & \infty & \infty & \infty & \infty & 1 & 0 \end{pmatrix}$$

**SVAR d:**

$$\begin{pmatrix} 0 & 1 & 2 & 2 & 3 & 3 & 4 & 5 \\ 1 & 0 & 1 & 1 & 2 & 2 & 3 & 4 \\ 2 & 1 & 0 & 2 & 3 & 3 & 4 & 5 \\ 2 & 1 & 2 & 0 & 1 & 1 & 2 & 3 \\ 3 & 2 & 3 & 1 & 0 & 2 & 3 & 4 \\ 3 & 2 & 3 & 1 & 2 & 0 & 1 & 2 \\ 4 & 3 & 4 & 2 & 3 & 1 & 0 & 1 \\ 5 & 4 & 5 & 3 & 4 & 2 & 1 & 0 \end{pmatrix}$$

**SVAR e:**

$$\begin{aligned} & (1+2+2+3+3+4+5)+(1+1+2+2+3+4)+(2+3+3+4+5)+(1+1+2+3)+(2+3+4)+(1+2)+1 \\ & = 20 + 13 + 17 + 7 + 9 + 3 + 1 = 70 \end{aligned}$$

**SVAR f:** 1-5, 1-6, 2-7, 3-5, 3-6, 4-8, 5-7. Total of 7.

**SVAR g:**

$$t_B = t_0 - \left( \frac{98}{n^2} (w_0 - \mathcal{W}(G)) + 5.5 \cdot (p_0 - p) \right)$$

$$t_0 = 745.42 \cdot \log_{10}(n + 4.4) - 689.4 = 745.42 \cdot \log_{10}(8 + 4.4) - 689.4 = 125.658$$

$$w_0 = \frac{1}{6} \cdot (n + 1) \cdot n \cdot (n - 1) = \frac{1}{6} \cdot (8 + 1) \cdot 8 \cdot (8 - 1) = 84$$

$$p_0 = n - 3 = 8 - 3 = 5$$

$$p = 7$$

$$t_B = 125.658 - \left( \frac{98}{8^2} (84 - 70) + 5.5 \cdot (5 - 7) \right) = 115.2205$$